



National Quail Symposium Proceedings

Volume 8

Article 19

2017

Effects of Broad-Scale Conservation on Northern Bobwhite Populations in Agricultural Landscapes

John M. Yeiser
University of Georgia

John J. Morgan
Kentucky Department of Fish and Wildlife Resources

Danna L. Baxley
Kentucky Department of Fish and Wildlife Resources

Richard B. Chandler
University of Georgia

James A. Martin
University of Georgia

Follow this and additional works at: <http://trace.tennessee.edu/nqsp>

 Part of the [Natural Resources and Conservation Commons](#)

Recommended Citation

Yeiser, John M.; Morgan, John J.; Baxley, Danna L.; Chandler, Richard B.; and Martin, James A. (2017) "Effects of Broad-Scale Conservation on Northern Bobwhite Populations in Agricultural Landscapes," *National Quail Symposium Proceedings*: Vol. 8 , Article 19.

Available at: <http://trace.tennessee.edu/nqsp/vol8/iss1/19>

This Plenary Session is brought to you for free and open access by Trace: Tennessee Research and Creative Exchange. It has been accepted for inclusion in National Quail Symposium Proceedings by an authorized editor of Trace: Tennessee Research and Creative Exchange. For more information, please contact trace@utk.edu.

EFFECTS OF BROAD-SCALE CONSERVATION ON NORTHERN BOBWHITE POPULATIONS IN AGRICULTURAL LANDSCAPES

John M. Yeiser¹

Warnell School of Forestry and Natural Resources, University of Georgia, 180 E Green St, Athens, GA 30602

John J. Morgan

Kentucky Department of Fish and Wildlife Resources, 1 Sportsman's Ln, Frankfort, KY 40603

Danna L. Baxley

Kentucky Department of Fish and Wildlife Resources, 1 Sportsman's Ln, Frankfort, KY 40603

Richard B. Chandler

Warnell School of Forestry and Natural Resources, University of Georgia, 180 E Green St, Athens, GA 30602

James A. Martin

Warnell School of Forestry and Natural Resources, University of Georgia, 180 E Green St, Athens, GA 30602

ABSTRACT

Private land initiatives such as the Conservation Reserve Enhancement Program (CREP) are avenues for broad-scale northern Bobwhite (*Colinus virginianus*) conservation. The CREP in Kentucky established 40,468 ha of native prairie grasses and riparian corridors in the Green River Basin. Northern bobwhite responses to similar conservation measures at local scales (i.e., the site of implementation) have been positive; however, the geographic extent of the influence of private land initiatives on populations is less understood. Our objectives were to investigate landscape-scale effects of CREP on northern bobwhite populations. Using a stratified random sampling design, 254 roadside point counts were performed over 5 years throughout the Green River Basin along a gradient of landscape-scale CREP density. Local-scale (500 m radius) CREP density was held constant at monitoring points. We analyzed data using an open-population distance sampling model that included estimators of appropriate landscape scale and strength of density dependence. Population response to the CREP was positive and outweighed conservation footprint. Our results suggest that broad-scale conservation can influence wildlife populations outside of targeted areas. Concurrently, because the majority of land in the Eastern U.S. is privately owned, private land conservation initiatives present an effective strategy for promoting wildlife population recovery across large areas. Our future directions with this research include improving model estimators, determining mechanisms behind landscape-scale effects of CREP, and determining the influence of the spatial arrangement of landscape features on local populations.

Citation: Yeiser, J. M., J. J. Morgan, D. L. Baxley, R. B. Chandler, and J. A. Martin. 2017. Effects of broad-scale conservation on northern bobwhite populations in agricultural landscapes. *National Quail Symposium Proceedings* 8:36.

Key words: northern bobwhite, Conservation Reserve Enhancement Program, broad-scale conservation, abundance estimation, private land initiatives

¹Email: john.yeiser@uga.edu